

General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

NASA TECHNICAL
MEMORANDUM

NASA TM X-73,204

NASA TM X-73,204

ANALYTICAL CALCULATION OF THE AREAS
OF SATURN'S DISK AND RINGS

S. Matthews and E. F. Erickson

Ames Research Center
Moffett Field, California 94035

February 1977



1. Report No. NASA TM X-73,204		2. Government Accession No.		3. Recipient's Catalog No.	
4. Title and Subtitle ANALYTICAL CALCULATION OF THE AREAS OF SATURN'S DISK AND RINGS				5. Report Date	
				6. Performing Organization Code	
7. Author(s) S. Matthews and E. F. Erickson				8. Performing Organization Report No. A-6906	
9. Performing Organization Name and Address NASA-Ames Research Center Moffett Field, Calif., 94035				10. Work Unit No. 352-03-03-02	
				11. Contract or Grant No.	
12. Sponsoring Agency Name and Address National Aeronautics and Space Administration Washington, D. C., 20546				13. Type of Report and Period Covered Technical Memorandum	
				14. Sponsoring Agency Code	
15. Supplementary Notes					
16. Abstract Determination of the thermal emission from the disk of Saturn at wavelengths in the vicinity of the thermal peak ($\sim 50 \mu\text{m}$) is complicated by the fact that infrared telescopes currently operating at these wavelengths cannot spatially separate the rings from the disk. To account for the emission from the rings, the area of the visible disk, the area of the ansae (visible rings not overlapping the disk), and the area of overlap (visible rings overlapping the disk) must be known. The calculation presented here describes the analytical determination of these areas from parameters available in the ephemeris.					
17. Key Words (Suggested by Author(s)) Astronomy			18. Distribution Statement Unlimited STAR Category - 89		
19. Security Classif. (of this report) Unclassified	20. Security Classif. (of this page) Unclassified		21. No. of Pages 6	22. Price* \$3.25	

ANALYTICAL CALCULATION OF THE AREAS

OF SATURN'S DISK AND RINGS

S. Matthews and E. F. Erickson

Ames Research Center

SUMMARY

Determination of the thermal emission from the disk of Saturn at wavelengths in the vicinity of the thermal peak ($\sim 50 \mu\text{m}$) is complicated by the fact that infrared telescopes currently operating at these wavelengths cannot spatially separate the rings from the disk. To account for the emission from the rings, the area of the visible disk, the area of the ansae (visible rings not overlapping the disk), and the area of overlap (visible rings overlapping the disk) must be known. The calculation presented here describes the analytical determination of these areas from parameters available in the ephemeris.

Saturn's rings and disk make comparable contributions to the thermally emitted flux from the entire system. If one assumes that the optical depth of the rings is not necessarily high, and that the only significant ring effects are due to the A and B rings, then five areas are important in estimating the relative contributions of the rings and disk. These are:

Ω_{vd} = unobscured (visible) area of the disk

Ω_A, Ω_B = visible area of the A and B rings respectively

ω_A, ω_B = area of disk obscured by the A and B rings respectively

The brightness of the entire system is described by brightness temperature T , which is related to the rings' temperatures, T_A and T_B , and the disk temperature, T_d , by

$$B_v(T)\Omega_t = B_v(T_d)(\Omega_{vd} + e^{-\tau_A}\omega_A + e^{-\tau_B}\omega_B) + B_v(T_A)(1 - e^{-\tau_A})\Omega_A + B_v(T_B)(1 - e^{-\tau_B})\Omega_B$$

Here B_v is the Planck function,

$$\Omega_t = \Omega_{vd} + \Omega_A + \Omega_B \quad (1)$$

is the total system area, and τ_A and τ_B are the optical depths of the rings.

Both the disk and rings are elliptical in appearance to the observer. In rectangular coordinates the equation of an ellipse is

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, \quad a \geq b \quad (2)$$

If we transform this to polar coordinates we have $x = r \cos \theta$, $y = r \sin \theta$, so

$$r^2 \left(\frac{\cos^2 \theta}{a^2} + \frac{\sin^2 \theta}{b^2} \right) = 1 \quad \text{or} \quad r(\theta) = ab(a^2 \sin^2 \theta + b^2 \cos^2 \theta)^{-1/2} \quad (3)$$

The area of an ellipse is πab . Our notation for the Saturn geometry is shown in figure 1. Thus, for the problem at hand, we have

$$\Omega_{vd} = \pi a_0 b_0 - \omega_A - \omega_B \quad (4)$$

$$\left. \begin{aligned} \Omega_A &= \pi(a_1 b_1 - a_2 b_2) - \omega_A \\ \Omega_B &= \pi(a_3 b_3 - a_4 b_4) - \omega_B \end{aligned} \right\} \quad (5)$$

If we note that (see ref. 1)

$$\frac{a_2}{a_1} = \frac{b_2}{b_1} = 0.8801 \quad (6)$$

$$\frac{a_3}{a_1} = \frac{b_3}{b_1} = 0.8599 \quad (7)$$

and

$$\frac{a_4}{a_1} = \frac{b_4}{b_1} = 0.6650 \quad (8)$$

then the quantities a_2 , b_2 , a_3 , b_3 , a_4 , and b_4 can be eliminated in (5) so that

$$\left. \begin{aligned} \Omega_A &= 0.2254 \pi a_1 b_1 - \omega_A \\ \Omega_B &= 0.2972 \pi a_1 b_1 - \omega_B \end{aligned} \right\} \quad (9)$$

a_0 , b_0 , a_1 , and b_1 can be obtained from the ephemeris.

There remains simply to calculate ω_A and ω_B . To determine the angles θ_i , $i = 1$ to 4, we note that

$$r_i(\theta_i) = r_0(\theta_i) \quad \text{or} \quad r_i^2(\theta_i) = r_0^2(\theta_i) \quad (10)$$

Then

$$\frac{a_1^2 b_1^2}{a_1^2 \sin^2 \theta_1 + b_1^2 \cos^2 \theta_1} = \frac{a_0^2 b_0^2}{a_0^2 \sin^2 \theta_1 + b_0^2 \cos^2 \theta_1} \quad (11)$$

which can be solved for $\tan \theta_1$

$$t_1 \equiv \tan \theta_1 = \frac{b_0 b_1}{a_0 a_1} \left(\frac{a_1^2 - a_0^2}{b_0^2 - b_1^2} \right)^{1/2} \quad \text{for } b_1 \leq b_0 \quad (12)$$

If b_1 should be greater than b_0 , $t_1 = +\infty$ [$\theta_1 = (\pi/2)$]. Now

$$\omega_A = 2 \int_{\theta_2}^{\theta_1} \int_{r_2}^{r_0} r \, dr \, d\theta + 2 \int_{\theta_1}^{\pi/2} \int_{r_2}^{r_1} r \, dr \, d\theta \quad (13)$$

$$= \int_{\theta_2}^{\theta_1} (r_0^2 - r_2^2) d\theta + \int_{\theta_1}^{\pi/2} (r_1^2 - r_2^2) d\theta \quad (14)$$

$$= \int_{\theta_2}^{\theta_1} r_0^2 d\theta + \int_{\theta_1}^{\pi/2} r_1^2 d\theta - \int_{\theta_2}^{\pi/2} r_2^2 d\theta \quad (15)$$

$$= a_0 b_0 \tan^{-1} \left(\frac{a_0}{b_0} \tan \theta \right) \Big|_{\theta_2}^{\theta_1} + a_1 b_1 \tan^{-1} \left(\frac{a_1}{b_1} \tan \theta \right) \Big|_{\theta_1}^{\pi/2} \\ - a_2 b_2 \tan^{-1} \left(\frac{a_2}{b_2} \tan \theta \right) \Big|_{\theta_2}^{\pi/2} \quad (16)$$

since

$$\int r^2 d\theta = a^2 b^2 \int \frac{d\theta}{a^2 \sin^2 \theta + b^2 \cos^2 \theta} = ab \tan^{-1} \left(\frac{a}{b} \tan \theta \right) \quad (17)$$

Rewriting (16), and from a similar expression for ω_B , we find

$$\omega_A = a_0 b_0 \left[\tan^{-1} \left(\frac{a_0}{b_0} t_1 \right) - \tan^{-1} \left(\frac{a_0}{b_0} t_2 \right) \right] + a_1 b_1 \left[\frac{\pi}{2} - \tan^{-1} \left(\frac{a_1}{b_1} t_1 \right) \right] \\ - a_2 b_2 \left[\frac{\pi}{2} - \tan^{-1} \left(\frac{a_2}{b_2} t_2 \right) \right] \quad (18)$$

and

$$\begin{aligned}\omega_B = a_0 b_0 \left[\tan^{-1} \left(\frac{a_0}{b_0} t_3 \right) - \tan^{-1} \left(\frac{a_0}{b_0} t_4 \right) \right] + a_3 b_3 \left[\frac{\pi}{2} - \tan^{-1} \left(\frac{a_3}{b_3} t_3 \right) \right] \\ - a_4 b_4 \left[\frac{\pi}{2} - \tan^{-1} \left(\frac{a_4}{b_4} t_4 \right) \right]\end{aligned}\quad (19)$$

From (6), (7), and (8) we obtain finally

$$\begin{aligned}\omega_A = a_0 b_0 \left[\tan^{-1} \left(\frac{a_0}{b_0} t_1 \right) - \tan^{-1} \left(\frac{a_0}{b_0} t_2 \right) \right] + a_1 b_1 \left[0.1127\pi - \tan^{-1} \left(\frac{a_1}{b_1} t_1 \right) \right. \\ \left. + 0.7746 \tan^{-1} \left(\frac{a_1}{b_1} t_2 \right) \right]\end{aligned}\quad (20)$$

and

$$\begin{aligned}\omega_B = a_0 b_0 \left[\tan^{-1} \left(\frac{a_0}{b_0} t_3 \right) - \tan^{-1} \left(\frac{a_0}{b_0} t_4 \right) \right] + a_1 b_1 \left[0.1486\pi - 0.7394 \tan^{-1} \left(\frac{a_1}{b_1} t_3 \right) \right. \\ \left. + 0.4422 \tan^{-1} \left(\frac{a_1}{b_1} t_4 \right) \right]\end{aligned}\quad (21)$$

As an example, we calculate the areas for January 27, 1976. On this date the ephemeris parameters (in arc seconds) are

$$\begin{aligned}a_0 &= 10.295 & b_0 &= 9.215 \\ a_1 &= 23.19 & b_1 &= 8.49\end{aligned}$$

Then, from (1), (4), (9), (20), and (21) one obtains in square arc seconds the areas

$$\begin{aligned}\Omega_{vd} &= 259.02 \\ \Omega_A &= 127.91 \\ \Omega_B &= 156.25 \\ \omega_A &= 11.52 \\ \omega_B &= 27.49 \\ \Omega_t &= 543.18\end{aligned}$$

REFERENCE

The American Ephemeris and Nautical Almanac, issued by the Nautical Almanac Office, United States Naval Observatory. U. S. Government Printing Office, Washington, 1976, p. 416.

SATURN GEOMETRY




Figure 1.- Notation for the calculation of the Saturn geometry. The quantities a_i and b_i are the semi-major and semi-minor axes of the indicated components, and the θ_i are the angles of intersection of the ring edges with the disk. The areas of overlap of the disk with ring A and ring B are ω_A and ω_B , respectively.